



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

varied in *terms* or *signs*. After the *Synthesis* our *Author* shews the *analysis* or *method* by which he found this Rule, viz. a *Parabola* being described, and a *point* in its *plain* given in *position*, he expresses 2 ways, the *radius* of a *Circle* passing through the *Vertex* of any *diameter*, i. e. by position of the given *Center*, and application of the foresaid *propriety* to express the *ratio* of the *radius* to the given lines of the *parabola*: So having an *Equation* of 4 *dimensions*, and rejecting equal on both sides, he depresses it to a *Cubic*, but adjoyning to it a quantity for the *Homogene* of the comparison, the *Equation* subsists in a *Biquadratic*, having all its *terms*, if the *Circle* be supposed to pass not thro the *vertex* of the *diameter*, but thro a *point* which being joyn'd with the *Vertex* and *Center* may terminate a right angled triangle.

This *Equation* he compares with another like it and equal to it; then by *equating* the *Coefficients* of these 2 *Equations* he presently discovers the *central* Rule; whose universal extent appears in *Biquadratic Equations* affected under all their *Parodic* degrees; for all the other *cases* where any *terms* are wanting, are but *Corollarys* or more compendious *Constructions* deriv'd from the general rule. So that the invention of the rule seems as much due to the last *Equation* of the *Coefficients*, as to the foresaid *propriety*, which is demonstrated by *Archimedes* in the *Section* of a *parabolic Conoid* by a *plane* parallel to the *axis*, and is particularly used by *Slusius* in his *Analytics*, who thereby constructs a *Biquadratic Equation* keeping all its *terms*. But then the *Analysis* of *Slusius* by breaking the *Equation* into 2 others to find 2 *places* is very different from that whereby our *Author* found his *central rule*; then which nothing can be expected more easie, simple, or universal; seeing any *Parabola* being once for all described, will give all the roots true and false, of any *Equation* without reduction or any alteration.

ERRATUM. p. 518. line the last, read Nubigenum.

O X F O R D, Printed at the THEATER, and are to be sold by *Moses Pitt*, at the *Angel*, and *Samuel Smith*, at the *Princes Arms* in *St. Paul's Church-yard* L O N D O N. 1684.